

STIC-ILL

TP248.3.77

Adm2

From: Helmer, Georgia  
Sent: Friday, October 18, 2002 11:37 AM  
To: STIC-ILL  
Subject: Satac-STN

COU YOU PLEASE GET THIS STUFF FOR MME? VERY MANY THANKS IN ADVANCE!!!!!!!!!!!!!!!!!!!!!!!!!!!!

Georgia L. Helmer Ph.D.  
Patent Examiner  
Crystal Mall 1, 9D14  
AU 1638  
703-308-7023  
Georgia.Helmer@USPTO.gov

AN 2002008965 MEDLINE  
DN 21237598 PubMed ID: 11338924  
TI Satellite DNA-based artificial chromosomes for use in gene therapy.  
AU Hadlaczky G  
CS Institute of Genetics, Biological Research Center, Hungarian Academy of  
Sciences, H-6701 Szeged, PO Box 521, Hungary.. hgy@nucleus.szbk.u-  
szeged.hu  
SO Curr Opin Mol Ther, (2001 Apr) 3 (2) 125-32. Ref: 33  
Journal code: 100891485. ISSN: 1464-8431.  
CY England: United Kingdom  
DT Journal; Article; (JOURNAL ARTICLE)  
General Review; (REVIEW)  
(REVIEW, TUTORIAL)  
LA English  
FS Priority Journals  
EM 200112  
ED Entered STN: 20020121  
Last Updated on STN: 20020121  
Entered Medline: 20011205

L7 ANSWER 2 OF 9 CAPLUS COPYRIGHT 2002 ACS  
D

ALL CITATIONS AVAILABLE IN THE RE FORMAT

AN 2000:716318 CAPLUS  
DN 134:232422  
TI Satellite DNA-based artificial chromosomes-chromosomal vectors. Reply to  
Comments  
AU Brown, William R. A.  
CS Institute of Genetics, University of Nottingham, Nottingham, UK  
SO Trends in Biotechnology (2000), 18(10), 403  
CODEN: TRBIDM; ISSN: 0167-7799  
PB Elsevier Science Ltd.  
DT Journal  
LA English

L7 ANSWER 5 OF 9 CAPLUS COPYRIGHT 2002 ACS

AN 2000:716317 CAPLUS  
DN 134:232421  
TI Satellite DNA-based artificial chromosomes-chromosomal vectors. Comments  
AU Perez, Carl; de Jong, Gary; Drayer, Jan  
CS Chromos Molecular Systems Inc., Burnaby, BC, Can.  
SO Trends in Biotechnology (2000), 18(10), 402-403  
CODEN: TRBIDM; ISSN: 0167-7799  
PB Elsevier Science Ltd.  
DT Journal  
LA English

L7 ANSWER 6 OF 9 MEDLINE DUPLICATE 2

AN 2000493348 MEDLINE  
DN 20297740 PubMed ID: 10841045  
TI Generation of transgenic mice and germline transmission of a mammalian

Enzymes show a delicate and complicated balance of all the factors that help determine the characteristics of the biocatalysts, and molecular modeling techniques can be extremely helpful in elucidating these interconnected factors. In the previously cited case of the thermostable mutant of subtilisin E, for example, the use of an all atom representation of the protein, its mutant and the solvent, together with the application of molecular dynamics to study the time evolution of the system, have allowed us to obtain a clear picture of enthalpic and entropic factors that determine enzyme stability and activity<sup>11</sup>. It was revealed that the penetration of solvent into the hydrophobic core of the native protein, which leads to unfolding, involves one region of the protein in particular. A semi-quantitative definition of flexibility is also given, helping to obtain a clear picture of the factors that determine the behaviour of subtilisin E and of its thermophilic homologue<sup>11</sup>.

Thus, the use of directed evolution (coupled with suitable screening methods) and molecular modeling techniques can become an extremely important factor in the elucidation

of the relationship between activity and stability in enzymes. The experimental results could be used by the modelers to increase their knowledge about the reasons of protein stability and to design better catalysts – enzymes that couple high activity and high stability – which would be a major achievement in the area of applied biocatalysis.

### Giacomo Carrea and Giorgio Colombo

Istituto di Biocatalisi e Riconoscimento Molecolare, CNR, Via Mario Bianco 9, 20131 Milano, Italy.  
(E-mail: gcarrea@ico.mi.cnr.it)

### References

- 1 Carrea, G. and Riva, S. (2000) Properties and synthetic applications of enzymes in organic solvents. *Angew. Chem. Int. Ed. Engl.* 39, 2226–2254
- 2 Hough, D.W. and Danson, M.J. (1999) Extremozymes. *Curr. Opin. Chem. Biol.* 3, 39–46
- 3 Zhang, T. *et al.* (1995) Enhancement of protein stability by the combination of point mutations in T4 lysozyme is additive. *Protein Eng.* 8, 1017–1022
- 4 Wrba, A. *et al.* (1990) Extremely thermostable D-glyceraldehyde-3-phosphate dehydrogenase from the eubacterium *Thermotoga maritima*. *Biochemistry* 29, 7584–7592
- 5 Varley, P.G. and Pain, R.H. (1991) Relation between stability, dynamics and enzyme activity in 3-phosphoglycerate kinases from yeast and *Thermus thermophilus*. *J. Mol. Biol.* 220, 531–538
- 6 Knapp, S. *et al.* (1997) Crystal structure of glutamate dehydrogenase from the hyperthermophilic eubacterium *Thermotoga maritima* at 3.0 Å resolution. *J. Mol. Biol.* 267, 916–932
- 7 Aguilar, C.F. *et al.* (1997) Crystal structure of the beta-glycosidase from the hyperthermophilic archaeon *Sulfolobus solfataricus*: resilience as a key factor in thermostability. *J. Mol. Biol.* 271, 789–802
- 8 Zhao, H. and Arnold, F.H. (1999) Directed evolution converts subtilisin E into a functional equivalent of thermolysin. *Protein Eng.* 12, 47–52
- 9 Zhao, H. *et al.* (1998) Molecular evolution by staggered extension process (StEP) in vitro recombination. *Nat. Biotechnol.* 16, 258–261
- 10 Lazaridis, T. *et al.* (1997) Dynamics and unfolding pathways of a hyperthermophilic and a mesophilic rubredoxin. *Protein Sci.* 6, 2589–2605
- 11 Colombo, G. and Merz, K.M. (1999) Stability and activity of mesophilic subtilisin E and its thermophilic homolog: insights from molecular dynamics simulations. *J. Am. Chem. Soc.* 121, 6895–6903

## Satellite DNA-based artificial chromosomes – chromosomal vectors

### Letter

**W**e were intrigued by a recent review<sup>1</sup> in which Brown *et al.* described two approaches to developing vectors that have the properties of chromosomes. We would like to bring attention to the existence of a third chromosome-based vector technology – satellite-DNA-based artificial chromosomes (SATACs; Refs 2,3) – which has been extensively reviewed<sup>4,5</sup>.

In the first protocol that Brown described – cellular-mediated chromosome assembly – chromosomes were assembled from transfected alphoid, telomeric and marker DNAs (Refs 6,7). The limitations of this approach are that: (1) there is no predictable relationship between input DNA and assembled chromosomes; (2) rearrangements frequently occur

when the transfected DNA fails to form neochromosomes; (3) the neochromosomes form at frequencies of  $\sim 5 \times 10^{-5}$ ; and (4) neochromosomes have been shown to form only in HT1080 human fibrosarcoma cells.

The second protocol involves fragmenting natural human chromosomes using telomere-directed breakage to generate minichromosomes<sup>8,9</sup>. Transgenic animals have been generated with human–murine minichromosome chimeras<sup>9</sup>, with fragmented human chromosomes<sup>10–12</sup>, and recently with human small accessory chromosomes (SACs) (P. Marynen *et al.*, unpublished data). The primary limitation of this protocol is the inefficient and tedious transfer of these minichromosomes from one cell type to another by microcell fusion.

A third approach was omitted from the review by Brown *et al.* This approach uses neochromosomes formed by the *de novo* amplification of pericentric heterochromatin, SATACs (Refs 2–4). Exogenous plasmid DNA is introduced into the vicinity of mouse<sup>2,3</sup> and human<sup>13</sup> pericentric heterochromatin, thus forming *de novo* dicentric chromosomes as the result of the amplification of endogenous pericentric heterochromatin and marker sequences. Ensuing breakage generates chromosomes that range from 10 to 360 megabases. The murine chromosomal vector has been transferred to hamster, human and bovine cells, and is transmitted as a discrete chromosome with segregation efficiencies per generation between 99.0% and 99.8% (Ref. 14). These chromosomes are largely composed of repetitive sequences, thus enabling high-throughput isolation using flow cytometry<sup>15</sup>. Transgenic mice have been generated using the unique process of microinjecting isolated SATACs into the pronuclei of fertilized murine oocytes<sup>16</sup>. This extra

chromosome has been transmitted through the germline over three generations and has been maintained as a discrete chromosome for over 14 months in lymphocytes (D.O. Co *et al.* unpublished data). Recently, transgenic founder mice and their progeny have been generated that carry a SATAC with a 1.0-megabase payload.

We have extended our studies on chromosome delivery beyond microcell fusion. SATACs isolated using flow cytometry have been transferred intact to other cell types by electroporation, microinjection and lipid-mediated techniques – generating stable transformants at efficiencies exceeding  $10^{-2}$  (de Jong *et al.*, unpublished data).

We believe that by using SATACs to deliver large DNA payloads in an efficient, safe and stable manner, we have taken the first steps in developing an artificial chromosome as an 'ideal vector'.

**Carl Perez  
Gary de Jong  
Jan Drayer**

*Chromos Molecular Systems Inc.,  
8081 Lougheed Highway, Burnaby,  
BC, Canada V5A 1W9.  
(E-mail: cperez@chromos.com)*

## Gyula Hadlaczky

*Institute of Genetics, Biological Research Center  
of the Hungarian Academy of Sciences,  
PO Box 521, H-6701, Szeged, Hungary.*

### References

- 1 Brown, W.R.A. *et al.* (2000) Artificial chromosomes: ideal vectors? *Trends Biotechnol.* 18, 218–223
- 2 Kereső, J. *et al.* (1996) *De novo* chromosome formations by large-scale amplification of the centromeric region of mouse chromosomes. *Chromosome Res.* 4, 226–239
- 3 Holló, G. *et al.* (1996) Evidence for a megareplicon covering megabases of centromeric chromosome segments. *Chromosome Res.* 4, 240–247
- 4 Vos, J.-M.H. (1998) Mammalian artificial chromosomes as tools for gene therapy. *Curr. Opin. Genet. Dev.* 8, 351–359
- 5 Stock, G. and Campbell, J. (2000) A vision for practical human germline engineering. In *Engineering the Human Germline: An Exploration of the Science and Ethics of Altering the Genes we pass to our Children* (Stock, G. and Campbell, J., eds), pp. 9–16. Oxford University Press
- 6 Harrington, J.J. *et al.* (1997) Formation of *de novo* centromeres and construction of first-generation human artificial microchromosomes. *Nat. Genet.* 15, 345–355
- 7 Ikeno, M. *et al.* (1998) Construction of YAC-based mammalian artificial chromosomes. *Nat. Biotechnol.* 16, 431–439
- 8 Shen, M.H. *et al.* (1997) Human mini-chromosomes in mouse embryonal stem cells. *Hum. Mol. Genet.* 6, 1375–1382
- 9 Shen, M.H. *et al.* (1999) A structurally defined mini-chromosome vector for the mouse germ line. *Curr. Biol.* 10, 31–34
- 10 Tomizuka, K. *et al.* (1997) Functional expression and germ line transmission of a human chromosome fragment in chimeric mice. *Nat. Genet.* 16, 133–143
- 11 Tomizuka, K. *et al.* (2000) Double trans-chromosomal mice: maintenance of two individual human chromosome fragments containing Ig heavy and k loci and expression of fully human antibodies. *Proc. Natl. Acad. Sci. U. S. A.* 97, 722–727
- 12 Hernandez, D. *et al.* (1999) Transchromosomal mouse embryonal stem cell lines and chimeric mice that contain freely segregating fragments of human chromosome 21. *Hum. Mol. Genet.* 8, 923–933
- 13 Csonka, E. *et al.* (2000) Human satellite DNA-based artificial chromosomes. *Eur. J. Hum. Genet. (Suppl.)* 8, 40
- 14 Telenius, H. *et al.* (1999) Stability of a functional murine satellite DNA-based artificial chromosome across mammalian species. *Chromosome Res.* 7, 3–7
- 15 de Jong, G. *et al.* (1999) Mammalian artificial chromosome pilot production facility: large scale isolation of functional satellite DNA-based artificial chromosomes. *Cytometry* 35, 129–133
- 16 Co, D.O. *et al.* (2000) Generation of transgenic mice and germline transmission of a mammalian artificial chromosome introduced into embryos by pronuclear microinjection. *Chromosome Res.* 8, 183–191

## Response from Brown

**W**e thank Perez *et al.* for their interest in our recent review and for providing the opportunity to discuss artificial chromosomes in more detail. In our review, we argued that artificial chromosomes are potentially interesting because they should enable some of the *dis-*acting sequences that are necessary for chromosome function to be identified and because they might enable long tracts of genomic DNA to be introduced into the genomes of livestock or crop plants. We argued that both of these uses need artificial chromosome vectors of defined structures and a technology that allows precise modification of the artificial chromosome molecules. Satellite-DNA-based artificial chromosomes (SATACs) do not have molecularly defined structures; we did not discuss SATACs in this review because we doubt that they

will ever be useful either for addressing questions of basic biological interest or for biotechnological purposes. Perez *et al.* have emphasized the potential value of their SATACs as vectors. However, the genes present on SATACs have not, to our knowledge, been identified. Thus, given the current difficulties in Europe and the US involving crop plants modified with a single gene, we find it hard to see how regulators will accept animals carrying a

chromosome that might bear many undefined genes. Perez *et al.* report unpublished data that describes efficient transfer of SATACs between cells. This point is potentially interesting because it bears upon a technical problem that is common to all the approaches towards artificial chromosome construction and we look forward to seeing the data in published form.

**William R.A. Brown**

*Institute of Genetics, University of Nottingham,  
Queen's Medical Centre, Nottingham,  
UK NG7 2UH.  
(E-mail: william.r.brown@nottingham.ac.uk)*

*Letter*

### Letters to the Editor

*Trends in Biotechnology* welcomes correspondence that are of interest to the biotechnology community. Please send letters to:

**TTEC@current-trends.com**